



Distorsal

1. Identification of the veterinary drug

WERAVET 13 Distorsal
Liquid dilution for injection

Active substance: Rhus Toxicodendron (HAB 34) C 30

For animals: horses, cattle, pigs, dogs

2. Qualitative and quantitative composition

1 ampoule of 2 ml (2010.1 mg) contains:

Medically active ingredient:

Rhus Toxicodendron (HAB 34) Dil. C 30 199.1 mg (HAB; V.2a)

Other components:

Sodium chloride
Water for injection

3. Pharmaceutical form

Liquid dilution for injection

4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)

Homeopathic drug

4.2 Pharmacokinetic characteristics

No specifications

4.3 Toxicological characteristics

Acute toxicity, subchronic and chronic toxicity

No examination results are available for the acute, subchronic and chronic toxicity of : Rhus Toxicodendron (HAB 34) C 30.

Mutagenic and tumour-causing potential

No examination results are available for the mutagenic and tumour-causing potential of : Rhus Toxicodendron (HAB 34) C 30.

Reproductive toxicity

No examination results are available for the reproductive toxicity of Rhus Toxicodendron (HAB 34) C 30.

5. Clinical specifications

5.1 Target animal species : horses, cattle, pigs, dogs

5.2 Indications :

The fields of application correspond to the veterinary-homeopathic drug pattern, and include:

- lameness in horses, cattle, pigs and dogs, (exacerbated by rest and improved by movement), frequently as a consequence of overexertions.
- acute, traumatic and chronic joint, ligament and tendon diseases as a consequence of:
 1. sprains
 2. torn ligaments
 3. tendonitis and peritendonitis

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- fever
- acute and strong inflammatory symptoms such as redness, heat, swelling, pain and dysfunction
- concurrent occurrence of swellings on several joints (polyarthritis)
- severe impairment of general well-being
- persistent, unclear, periodic or emergent symptoms
- deficiency states resulting from reduced intake of vital nutrients

The use of WERAVET 13 Distorsal should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 13 Distorsal may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life
- disease which require surgical treatment, such as ruptures of tendons, muscles or ligaments and fractures.

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

5.3 Contraindications

None known

5.4 Side effects (type, frequency and degree of severity)

Note:

In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 13 Distorsal during gestation and lactation.

5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 13 Distorsal should be avoided. Please consult your veterinarian.

5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection.

horses	3-4 ml
cattle	3-4 ml
pigs	2-3 ml
dogs	1-2 ml

In acute conditions, the injection should be repeated after 1 – 2 days; otherwise, repetition takes place after 4 – 8 days.

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.



Distorsal

5.9 Overdose (symptoms, emergency measures and antidote)

No specifications

5.10 Special warnings for every target animal species

No specifications

5.11 Waiting period

Horses edible tissues: 0 days
milk: 0 days

Cattle: edible tissues: 0 days
milk: 0 days

Pigs edible tissues: 0 days

5.12 Special precautionary measures for the user

No specifications

6. Pharmaceutical specifications

6.1 Incompatibilities

No specifications

6.2 Duration of shelf life

6.2.1 of the finished drug in the undamaged container
5 years

6.2.2 of the finished drug after the container is opened

Use immediately after opening. Opened ampoules must be discarded.

6.2.3 after mixing the preparation, ready for use

Not applicable

6.3 Special notes regarding storage

Do not store above 25 °C.

6.4 Type and content of the container (package sizes)

Original packaging with 10 and 40 ampoules à 2 ml (clear glass)

6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment

Not applicable

7. Name or company and address of the pharmaceutical entrepreneur

Dr. Assmann
Veterinärspécialitäten GmbH
Kehler Str. 7
76437 Rastatt
Germany
www.weravet.de

Also distributed by:

Biokanol Pharma GmbH
Kehler Str. 7
76437 Rastatt
Germany
Tel.: +49 7222-78679-0
Fax: +49 7222-78679-9
www.biokanol.de

8. License number
6157753.00.00

9. Date of licensing / renewal
26.03.2004

10. Prescription status / prescription drug in Germany:
non prescription drug

11. Date of information
March 2004