



Dysenteral

1. Identification of the veterinary drug

WERAVET 8 Dysenteral
Liquid dilution for injection

For animals: : horses, cattle, pigs, sheep, goats, dogs, cats

2. Qualitative and quantitative composition

1 ampoule of 2 ml (2010.1 mg) contains:

Medically active ingredient:

Acidum arsenicosum Dil. C 30 199,1 mg
Podophyllum peltatum Dil. C 30 199,1 mg
Rheum Dil. C 30 mg 199,1 mg

Other components:

Sodium Chloride
Water for injection

3. Pharmaceutical form

Liquid dilution for injection

4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)
Homeopathic drug

4.2 Pharmacokinetic characteristics
No specifications

4.3 Toxicological characteristics
Acute toxicity, subchronic and chronic toxicity
No examination results are available for the acute, subchronic and chronic toxicity of
Acidum arsenicosum Dil. C 30,
Podophyllum peltatum Dil. C 30,
Rheum Dil. C 30 mg

Mutagenic and tumour-causing potential
No examination results are available for the mutagenic and tumour-causing potential of
Acidum arsenicosum Dil. C 30,
Podophyllum peltatum Dil. C 30,
Rheum Dil. C 30

Reproductive toxicity
No examination results are available for the reproductive toxicity of Acidum arsenicosum Dil. C 30,
Podophyllum peltatum Dil. C 30,
Rheum Dil. C 30

5. Clinical specifications

5.1 Target animal species:
horses, cattle, pigs, sheep, goats, dogs, cats

5.2 Indications
The fields of application correspond to the veterinary-homeopathic drug pattern, and include:

- diarrheal illnesses in the specified animal species

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- diarrheal illnesses lasting more than 2 days
- cases of diarrhea accompanied by admixtures of blood or black coloration of the stool
- high fever or fever that lasts more than 3 days accompanying circulatory insufficiency
- states of agitation
- severe impairment of general well-being
- acute and strong inflammatory symptoms such as redness, heat, swelling, pain and dysfunction.
- lasting, unclear, periodic or newly appearing complaints

In case of severe or persistent diarrhoeas the veterinary surgeon is to decide on the necessity of a continuing treatment.

The use of WERAVET 8 Dysenteral should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 8 Dysenteral may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life
- disease which require surgical treatment.

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

5.3 Contraindications
None known

5.4 Side effects (type, frequency and degree of severity)

Note:
In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 8 Dysenteral during gestation and lactation.

Like all drugs, homeopathic drugs should only be used during gestation and lactation after consultation with the veterinarian.

5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 8 Dysenteral should be avoided. Please consult your veterinarian.

5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection.

horses	3-4 ml
cattle	3-4 ml
pigs	2-3 ml
sheep	2-3 ml
goats	2-3 ml
dogs	1-2 ml
cats	1-2 ml

Dysenteral

In acute conditions, the injection should be repeated after 1 – 2 days; afterwards treatment should be continued with Dysenteral drops until clinical symptoms have disappeared completely

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.

5.9 Overdose (symptoms, emergency measures and antidote)

No specifications

5.10 Special warnings for every target animal species

No specifications

5.11 Waiting period

horses	edible tissues:	0 days
	milk:	0 days
cattle:	edible tissues:	0 days
	milk:	0 days
sheep	edible tissues:	0 days
	milk:	0 days
goats	edible tissues:	0 days
	milk:	0 days
pigs	edible tissues:	0 days

5.12 Special precautionary measures for the user

No specifications

6. Pharmaceutical specifications

6.1 Incompatibilities

No specifications

6.2 Duration of shelf life

6.2.1 of the finished drug in the undamaged container

5 years

6.2.2 of the finished drug after the container is opened

Use immediately after opening. Opened ampoules must be discarded.

6.2.3 after mixing the preparation, ready for use

Not applicable

6.3 Special notes regarding storage

Do not store above 25 °C.

6.4 Type and content of the container (package sizes)

Original packaging with 20 ampoules à 2 ml (clear glass)

6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment

Not applicable

7. Name or company and address of the pharmaceutical entrepreneur

Dr. Assmann
Veterinärspécialitäten GmbH
Kehler Str. 7
76437 Rastatt
Germany
www.weravet.de

Also distributed by:

Biokanol Pharma GmbH
Kehler Str. 7
76437 Rastatt
Germany
Tel.: +49 7222-78679-0
Fax: +49 7222-78679-9
www.biokanol.de

8. License number

6158451.00.00

9. Date of licensing / renewal

06.06.2003

10. Prescription status / prescription drug in Germany

Non prescription drug

11. Date of information

July 2003