



## Sangostyptal

### 1. Identification of the veterinary drug

WERAVET 16 Sangostyptal  
Liquid dilution for injection

For animals: : horses, cattle, pigs, sheep, goats, dogs, cats

### 2. Qualitative and quantitative composition

1 ampoule of 2 ml (2010.1 mg) contains:  
Medically active ingredient:  
Hamamelis virginiana Dil. C 30 199,1 mg  
Achillea millefolium Dil. C 30 199,1 mg

#### Other components:

Sodium Chloride  
Water for injection

### 3. Pharmaceutical form

Liquid dilution for injection

### 4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

**4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)**  
Homeopathic drug

**4.2 Pharmacokinetic characteristics**  
No specifications

**4.3 Toxicological characteristics**  
*Acute toxicity, subchronic and chronic toxicity*  
No examination results are available for the acute, subchronic and chronic toxicity of Hamamelis virginiana Dil. C 30 and Achillea millefolium Dil. C 30

*Mutagenic and tumour-causing potential*  
No examination results are available for the mutagenic and tumour-causing potential of Hamamelis virginiana Dil. C 30 and Achillea millefolium Dil. C 30

*Reproductive toxicity*  
No examination results are available for the reproductive toxicity of Hamamelis virginiana Dil. C 30 and Achillea millefolium Dil. C 30

### 5. Clinical specifications

**5.1 Target animal species**  
horses, cattle, pigs, sheep, goats, dogs, cats

### 5.2 indications

The fields of application correspond to the veterinary-homeopathic drug pattern, and include:

- haemostatic agent for venous and arterial haemorrhages
- blood in milk

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- severe and persistent haemorrhages
- pulsing arterial (bright red) haemorrhages
- hereditary or acquired coagulation defects
- accompanying circulatory insufficiency
- severe impairment of general well-being
- pale mucous membranes, paleness and coldness of the affected regions
- intense reddening of the wound margins
- discolorations of the surrounding skin
- fever
- acute and strong signs of inflammation, such as redness, heat, swelling, pain and function disturbance
- lasting, unclear, periodic or newly appearing complaints

In case of severe and persistent haemorrhages or haemorrhages after severe trauma the veterinary surgeon is to decide of the necessity of a continuing treatment.

The use of WERAVET 16 Sangostyptal should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 16 Sangostyptal may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life

- illnesses which require surgical treatment e.g. haemorrhages for unknown reasons, after trauma of extensive and deep injuries of the skin.

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

### 5.3 Contraindications

None known

### 5.4 Side effects (type, frequency and degree of severity)

**Note:**  
In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

### 5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

### 5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 16 Sangostyptal during gestation and lactation.

Like all drugs, homeopathic drugs should only be used during gestation and lactation after consultation with the veterinarian.

### 5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 16 Sangostyptal should be avoided. Please consult your veterinarian.



## Sangostyptal

### 5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection.

|        |        |
|--------|--------|
| horses | 3-4 ml |
| cattle | 3-4 ml |
| pigs   | 2-3 ml |
| sheep  | 2-3 ml |
| goats  | 2-3 ml |
| dogs   | 1-2 ml |
| cats   | 1-2 ml |

In acute conditions, the injection should be repeated after 1 – 2 days; otherwise, repetition takes place after 4 – 8 days. In peracute cases the single dose for stopping bleeding arterial and venous haemorrhages may be repeated in intervals of 5-10 minutes

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.

### 5.9 Overdose (symptoms, emergency measures and antidote)

No specifications

### 5.10 Special warnings for every target animal species

No specifications

### 5.11 Waiting period

|         |                 |        |
|---------|-----------------|--------|
| Horses  | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Cattle: | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Sheep   | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Goats   | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Pigs    | edible tissues: | 0 days |

### 5.12 Special precautionary measures for the user

No specifications

## 6. Pharmaceutical specifications

### 6.1 Incompatibilities

No specifications

### 6.2 Duration of shelf life

#### 6.2.1 of the finished drug in the undamaged container

5 years

#### 6.2.2 of the finished drug after the container is opened

Use immediately after opening. Opened ampoules must be discarded.

#### 6.2.3 after mixing the preparation, ready for use

Not applicable

### 6.3 Special notes regarding storage

Do not store above 25 °C.

#### 6.4 Type and content of the container (package sizes)

Original packaging with 10 and 40 ampoules à 2 ml (clear glass)

#### 6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment

Not applicable

## 7. Name or company and address of the pharmaceutical entrepreneur

Dr. Assmann  
Veterinärspécialitäten GmbH  
Kehler Str. 7  
76437 Rastatt  
[www.weravet.de](http://www.weravet.de)

#### Also distributed by:

Biokanol Pharma GmbH  
Kehler Str. 7  
76437 Rastatt  
Tel.: +49 7222-78679-0  
Fax: +49 7222-78679-9  
[www.biokanol.de](http://www.biokanol.de)

### 8. License number

6158149.00.00

### 9. Date of licensing / renewal

05.05.2003

### 10. Prescription status / prescription drug in Germany

Non prescription drug

### 11. Date of the information

July 2003