



Staphylosal

1. Identification of the veterinary drug

WERAVET 10 Staphylosal
Liquid dilution for injection

Active substance: Hepar sulfuris C 30

For animals: : horses, cattle, pigs, sheep, goats, dogs, cats

2. Qualitative and quantitative composition

1 ampoule of 2 ml (2010.1 mg) contains:

Medically active ingredient:

Hepar sulfuris Dil. C 30 199,1 mg

Other components:

Sodium Chloride

Water for injection

3. Pharmaceutical form

Liquid dilution for injection

4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)

Homeopathic drug

4.2 Pharmacokinetic characteristics

No specifications

4.3 Toxicological characteristics

Acute toxicity, subchronic and chronic toxicity

No examination results are available for the acute, subchronic and chronic toxicity of Hepar sulfuris C 30.

Mutagenic and tumour-causing potential

No examination results are available for the mutagenic and tumour-causing potential of Hepar sulfuris C 30.

Reproductive toxicity

No examination results are available for the reproductive toxicity of Hepar sulfuris C 30.

5. Clinical specifications

5.1 Target animal species

horses, cattle, Pigs, sheep, goats, dogs, cats

5.2 indications

The fields of application correspond to the veterinary-homeopathic drug pattern, and include:

- all types of suppuration
- navel infections
- inveterate cases of bronchitis in large animals
- bilical infection
- abscess formation
- protracted bronchial infection in large animals
- otitis externa in dogs

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- fever
- acute and strong signs of inflammation, such as redness, heat, swelling, pain and function disturbance
- severe impairment of general well-being
- restlessness
- haemorrhages
- purulent or bloody sputum
- long lasting skin alterations
- discolorations of the surrounding skin and intense reddening of the wound margins
- lasting, unclear, periodic or newly appearing complaints

In case of the clinical pictures "abscess formation", "protracted bronchial infection in large animals" and in clinical pictures along with an injury or severe alterations of the skin surface the veterinary surgeon is to decide on the necessity of an antibiotic treatment

The use of WERAVET 10 Staphylosal should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 10 Staphylosal may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life
- illnesses which require surgical treatment as

extensive injuries, inner haemorrhages, diffuse phlegmones, foreign body abscess and abscess forming fistula in body caves

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

5.3 Contraindications

None known

5.4 Side effects (type, frequency and degree of severity)

Note:

In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 10 Staphylosal during gestation and lactation. Like all drugs, homeopathic drugs should only be used during gestation and lactation after consultation with the veterinarian.

5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 10 Staphylosal should be avoided. Please consult your veterinarian.

5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection.

horses	3-4 ml
cattle	3-4 ml
pigs	2-3 ml
sheep	2-3 ml
goats	2-3 ml
dogs	1-3 ml
cats	1-3 ml

Staphylosal

In acute conditions, the injection should be repeated after 1 – 2 days; otherwise, repetition takes place after 4 – 8 days.

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.

5.9 Overdose (symptoms, emergency measures and antidote)

No specifications

5.10 Special warnings for every target animal species

No specifications

5.11 Waiting period

Horses	edible tissues:	0 days
	milk:	0 days
Cattle:	edible tissues:	0 days
	milk:	0 days
Sheep	edible tissues:	0 days
	milk:	0 days
Goats	edible tissues:	0 days
	milk:	0 days
Pigs	edible tissues:	0 days

5.12 Special precautionary measures for the user

No specifications

6. Pharmaceutical specifications

6.1 Incompatibilities

No specifications

6.2 Duration of shelf life

6.2.1 of the finished drug in the undamaged container

5 years

6.2.2 of the finished drug after the container is opened

Use immediately after opening. Opened ampoules must be discarded.

6.2.3 after mixing the preparation, ready for use

Not applicable

6.3 Special notes regarding storage

Do not store above 25 °C.

6.4 Type and content of the container (package sizes)

Original packaging with 10 and 40 ampoules à 2 ml (clear glass)

6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment

Not applicable

7. Name or company and address of the pharmaceutical entrepreneur

Dr. Assmann
Veterinärspécialitäten GmbH
Kehler Str. 7
76437 Rastatt
www.weravet.de

Also distributed by:

Biokanol Pharma GmbH
Kehler Str. 7
76437 Rastatt
Tel.: +49 7222-78679-0
Fax: +49 7222-78679-9
www.biokanol.de

8. License number

6157948.00.00

9. Date of licensing / renewal

20.12.2002

10. Prescription status / prescription drug in Germany

Non prescription drug

11. Date of the information

June 2003