



## Synosal

### 1. Identification of the veterinary drug

WERAVET 9 Synosal  
Liquid dilution for injection

Active Substance :  
Acidum benzoicum e resina C 30

For animals: : horses, cattle, pigs, dogs, chicken, ornament birds

### 2. Qualitative and quantitative composition

1 ampoule of 2 ml (2010.1 mg) contains:

#### Medically active ingredient:

Acidum benzoicum e resina Dil. C 30  
199,1 mg

#### Other components:

Sodium Chloride  
Water for injection

### 3. Pharmaceutical form

Liquid dilution for injection

### 4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

#### 4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)

Homeopathic drug

#### 4.2 Pharmacokinetic characteristics

No specifications

#### 4.3 Toxicological characteristics

##### *Acute toxicity, subchronic and chronic toxicity*

No examination results are available for the acute, subchronic and chronic toxicity of Acidum benzoicum e resina C 30.

##### *Mutagenic and tumour-causing potential*

No examination results are available for the mutagenic and tumour-causing potential of Acidum benzoicum e resina C 30.

##### *Reproductive toxicity*

No examination results are available for the reproductive toxicity of Acidum benzoicum e resina C 30.

### 5. Clinical specifications

#### 5.1 Target animal species

horses, cattle, pigs, dogs, chicken, ornament birds

### 5.2 Indications

The fields of application correspond to the veterinary-homeopathic drug pattern, and include:

- disorders of the joints and urinary tract as a consequence of infections or from disturbances of uric acid metabolism

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- blood in urine (hematuria)
- urine excretion disorders
- acute joint complaints and instances of lameness
- fever
- acute and strong signs of inflammation, such as redness, heat, swelling, pain and function disturbance
- severe reduction in general state
- lasting, unclear, periodic or newly appearing complaints

In case of the clinical picture "disorders of the joints and urinary tract as a consequence of infections or from disturbances of uric acid metabolism" the veterinary surgeon is to decide on the necessity of an allopathic or antibiotic treatment.

The use of WERAVET 9 Synosal should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 9 Synosal may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life
- illnesses which require surgical treatment.

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

### 5.3 Contraindications

None known

### 5.4 Side effects (type, frequency and degree of severity)

Note:

In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

### 5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

### 5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 9 Synosal during gestation and lactation.

Like all drugs, homeopathic drugs should only be used during gestation and lactation after consultation with the veterinarian.

### 5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 9 Synosal should be avoided. Please consult your veterinarian.

### 5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection.

|                |          |
|----------------|----------|
| horses         | 3-4 ml   |
| cattle         | 3-4 ml   |
| pigs           | 2-3 ml   |
| dogs           | 1-2 ml   |
| chicken        | 0,5-1 ml |
| ornament birds | 0,5-1 ml |

## Synosal

In acute conditions, the injection should be repeated after 1 – 2 days; otherwise, repetition takes place after 4 – 8 days.

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.

**5.9 Overdose (symptoms, emergency measures and antidote)**  
No specifications

**5.10 Special warnings for every target animal species**  
No specifications

**5.11 Waiting period**

|         |                 |        |
|---------|-----------------|--------|
| Horses  | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Cattle: | edible tissues: | 0 days |
|         | milk:           | 0 days |
| Pigs    | edible tissues: | 0 days |
| Chicken | edible tissues: | 0 days |
|         | Eggs            | 0 days |

**5.12 Special precautionary measures for the user**  
No specifications

**6. Pharmaceutical specifications**

**6.1 Incompatibilities**  
No specifications

**6.2 Duration of shelf life**

**6.2.1 of the finished drug in the undamaged container**  
5 years

**6.2.2 of the finished drug after the container is opened**  
Use immediately after opening. Opened ampoules must be discarded.

**6.2.3 after mixing the preparation, ready for use**  
Not applicable

**6.3 Special notes regarding storage**  
Do not store above 25 °C.

**6.4 Type and content of the container (package sizes)**  
Original packaging with 10 ampoules à 2 ml (clear glass)

**6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment**  
Not applicable

**7. Name or company and address of the pharmaceutical entrepreneur**

Dr. Assmann  
Veterinärspécialitäten GmbH  
Kehler Str. 7  
76437 Rastatt  
[www.weravet.de](http://www.weravet.de)

Also distributed by:

Biokanol Pharma GmbH  
Kehler Str. 7  
76437 Rastatt  
Tel.: +49 7222-78679-0  
Fax: +49 7222-78679-9  
[www.biokanol.de](http://www.biokanol.de)

**8. License number**  
6157902.00.00

**9. Date of licensing / renewal**  
01.03.2004

**10. Prescription status / prescription drug in Germany**  
Non prescription drug

**11. Date of the information**  
March 2004