

Alleosal

1. Identification of the veterinary drug

WERAVET 20 Alleosal
Liquid dilution for injection

For animals: : zoological animals, dogs and cats, small animals

2. Qualitative and quantitative composition

1 ampoule of 2 ml (2 g) contains:

Medically active ingredients:

Apis mellifica C6 200 mg,
Arsenum jodatum C12 200mg,
Urtica urens C6 200mg

Other components:

Sodium Chloride
Water for injection

3. Pharmaceutical form

Liquid dilution for injection

4. Pharmacological and toxicological characteristics and, insofar as it serves the purpose for therapeutic use, specifications on pharmacokinetics

4.1 Pharmacodynamic characteristics (including specifications on the substance or indication group; if applicable, ATC vet. Code)

Homeopathic drug

4.2 Pharmacokinetic characteristics

No specifications

4.3 Toxicological characteristics

Acute toxicity, subchronic and chronic toxicity

No examination results are available for the acute, subchronic and chronic toxicity of

Apis mellifica C6,
Arsenum jodatum C12 or
Urtica urens C6

Mutagenic and tumour-causing potential

No examination results are available for the mutagenic and tumour-causing potential of

Apis mellifica C6,
Arsenum jodatum C12 or
Urtica urens C6

Reproductive toxicity

No examination results are available for the reproductive toxicity of Aristolochia

Apis mellifica C6,
Arsenum jodatum C12 or
Urtica urens C6

5. Clinical specifications

5.1 Target animal species :

zoological animals, dogs and cats, small animals

5.2 Indications

Registered homeopathic drug, therefore without naming a therapeutic indication

Use of the drug should not take place without the advice of a veterinarian and does not replace other drugs and measures prescribed by the veterinarian in this regard.

If the following signs appear, you should consult the veterinarian, since an illness which requires a veterinarian's diagnosis may be present:

- fever
- acute and strong inflammatory symptoms such as redness, heat, swelling, pain and dysfunction
- severe impairment of general well-being
- lasting, unclear, periodic or newly appearing complaints

The use of WERAVET 20 Alleosal should take place with consideration of the homeopathic "simile principle". For this purpose, the consultation of a veterinarian for the purpose of determining the disease pattern (homeopathic anamnesis) is recommended.

In the following illnesses, WERAVET 20 Alleosal may be used only as supporting treatment:

- conditions of insufficiency due to reduced absorption or availability of nutrients required for life
- disorders which require surgical treatment

Prior to the commencement of treatment, the causes of the complaints should therefore be largely clarified.

5.3 Contraindications

None known

5.4 Side effects (type, frequency and degree of severity)

Note:

In treatment with a homeopathic drug, the existing complaints may temporarily worsen (initial worsening). In this case, you should terminate the use of the drug and consult the veterinarian.

5.5 Special precautionary measures for use

Due to the risk of a parenteral application of a drug, the administration of a drug with the same method of acting which is licensed for oral or rectal application is to be preferred.

Indications for intravenous injections are:

- immediate occurrence of the effect
- local irritations of the tissues in subcutaneous or intramuscular injection, due to the components and/or amount of the preparation.

5.6 Use during gestation and lactation

No examination results are available regarding the safety of using WERAVET 20 Alleosal during gestation and lactation.

Like all drugs, homeopathic drugs should only be used during gestation and lactation after consultation with the veterinarian.

5.7 Interactions with other drugs and other interactions

The simultaneous administration of an incompatible drug which may negate the effect of WERAVET 20 Alleosal should be avoided. Please consult your veterinarian.

5.8 Dosing with individual and daily specifications, type and duration of use

For intravenous, subcutaneous and intramuscular injection:

Zoological animals	1-3 ml
dogs	1-2 ml
cats	1-2 ml
small animals, not being used as food animals, depending on their size	1-2-ml

One injection for 2-3 days, repeat depending on the condition of the animal.

Administration of the drug should not take place longer than to the complete healing of the animal: A homeopathic drug can trigger a disease pattern which corresponds to the homeopathic drug pattern (symptomatics) in healthy animals.



Alleosal

5.9 Overdose (symptoms, emergency measures and antidote)

No specifications

5.10 Special warnings for every target animal species

No specifications

5.11 Waiting period

Not applicable

Not to be used in animals classed as food animals

5.12 Special precautionary measures for the user

No specifications

6. Pharmaceutical specifications

6.1 Incompatibilities

No specifications

6.2 Duration of shelf life

6.2.1 of the finished drug in the undamaged container

5 years

6.2.2 of the finished drug after the container is opened

Use immediately after opening. Opened ampoules must be discarded.

6.2.3 after mixing the preparation, ready for use

Not applicable

6.3 Special notes regarding storage

Do not store above 25 °C.

6.4 Type and content of the container (package sizes)

Original packaging with 10, 40 and 100 ampoules à 2 ml (clear glass)

6.5 Special precautionary measures for the disposal of unused drugs or other special precautionary measures to avoid dangers to the environment

Not applicable

7. Name or company and address of the pharmaceutical entrepreneur

Dr. Assmann
Veterinärspécialitäten GmbH
Kehler Str. 7
76437 Rastatt
Germany
www.weravet.de

Also distributed by:

Biokanol Pharma GmbH
Kehler Str. 7
76437 Rastatt
Germany
Tel.: +49 7222-78679-0
Fax: +49 7222-78679-9
www.biokanol.de

8. License number

400444.00.00

9. Date of licensing / renewal

08.03.2001

10. Prescription status / prescription drug in Germany

non prescription drug

11. Date of the information

March 2001